



The ABC Model of Flower Development: Genetic Blueprint of Floral Architecture

Neelesh Vimal¹
Prashant Gigaulia²
Keerti Tantwai³
Yogendra Singh^{4*}

¹Ph.D. Scholar, Biotechnology
Centre, JNKVV, Jabalpur

²Guest Faculty, Biotechnology
Centre, JNKVV, Jabalpur

³Director, Biotechnology Centre
JNKVV, Jabalpur

⁴Assistant Professor
(Senior Scale) Biotechnology
Department of Genetics and
Plant Breeding, JNKVV,
Jabalpur – 482004 (M.P.)



Open Access

*Corresponding Author
Yogendra Singh*

Article History

Received: 14.08.2026

Revised: 19.08.2026

Accepted: 24.08.2026

This article is published under the
terms of the [Creative Commons
Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/).

INTRODUCTION

Angiosperms, or flowering plants, signify the most evolutionarily progressed and ecologically prevailing classification in the plant kingdom. Their remarkable success in terrestrial ecosystems is primarily due to the development of highly specialized reproductive organs called flowers. Apart from their visual appeal, flowers serve as intricate reproductive structures that orchestrate gamete generation, pollination, fertilization, and later seed and fruit formation. The massive variety seen in floral structures among angiosperm species signifies a complex network of genetic and molecular regulatory systems that accurately govern floral organ identity, arrangement, and growth. A significant breakthrough in plant developmental biology was the creation of the ABC model of flower development, an influential genetic model that clarifies how particular combinations of homeotic genes dictate the identity of floral organs in different concentric whorls of a flower. In 1988 George Haughn and Chris Somerville introduced this model concept, the ABC model transformed the comprehension of floral morphogenesis by showing that a limited number of regulatory genes can produce the complete pattern of floral organs.

This model proposes that the identity of floral organs is determined by the combined functions of three primary classes of homeotic genes A, B, and C genes. The ABC model originated mainly from comprehensive molecular genetic research carried out in the model plants *Arabidopsis thaliana* and *Antirrhinum majus*. These genes control the development of four floral whorls: sepals, petals, stamens, and carpels. Class A genes merely determine sepals in the first whorl, the joint action of A and B genes governs petals in the second whorl, B and C genes together formed stamens in the third whorl, and Class C genes independently regulate carpel development in the fourth whorl.

The ABC model recognised a molecular basis for comprehending floral organogenesis and concrete the way for more complex ideas like the ABCDE and quartet models of flower development. Presently, it stands as one of the most significant theories in plant genetics and evolutionary developmental biology, providing essential knowledge regarding floral evolution, agricultural enhancement, reproductive biology, and the genetic modification of flowering plants.

Basic Structure of a Flower

A standard dicot flower consists of four concentric whorls of organs arranged from the outermost to innermost positions:


1. **Whorl 1 – Sepals:** green, protective organs forming the calyx.
2. **Whorl 2 – Petals:** often colored and fragrant, forming the corolla.
3. **Whorl 3 – Stamens:** male reproductive organs composed of a filament and anther (androecium).

4. **Whorl 4 – Carpels:** female reproductive organs forming the gynoecium, containing ovules.

The specification of each whorl involves a unique combination of transcriptional regulators that instruct floral meristem cells to adopt particular organ identities. These regulators are encoded by the so-called floral homeotic genes, whose loss or misexpression leads to characteristic organ transformations (homeotic mutations).

Exposure Floral Homeotic Mutants

In *Arabidopsis* and *Antirrhinum*, traditional genetic screenings uncovered mutants where one floral organ type was substituted for alternative. For an example, certain mutants generated petals in place of sepals, or carpels rather than stamens. These mutants were crucial in unravelling the major genetic principles of floral organ identity.



In Arabidopsis, three main types of mutants were discovered:

A) Class A mutants (such as, *apetala1*, *apetala2*)

Sepals changed into carpels, and petals substituted by stamens.

B) Class B mutations (for example, *apetala3*, *pistillata*)

Petals changed into sepals, and stamens turned into carpels.

C) Class C mutations (e.g., *agamous*)

Stamens became petals, and another flower took the place of carpels.

These outcomes prompted the hypothesis that the identity of organs in each whorl arises from the combinatorial activity of A, B, and C gene functions, with these genes functioning in mutually antagonistic patterns within the floral meristem.

The Classical ABC Model

The ABC model proposes that three classes of homeotic gene functions determine the identity of floral organs in specific whorls through combinatorial expression:

Whorl	Organ Type	Gene Functions Active
1	Sepal	A
2	Petal	A+B
3	Stamen	B+C
4	Carpel	C

- **A-function genes** specify sepals and petals (when combined with B).
- **B-function genes** specify petals (with A) and stamens (with C).
- **C-function genes** specify stamens and carpels and also terminate floral meristem activity.

A crucial element of the model is that A and C functions mutually suppress each other, with the existence of one inhibiting the expression of the other in shared areas. This spatial limitation guarantees the correct formation of organ identity regions.

Molecular Basis of the ABC Genes

The majority of ABC genes code for MADS-box transcription factors, which are a group of DNA-binding proteins that control gene expression in the process of floral organ development. The MADS domain is a conserved motif for DNA binding (~58 amino acids) designated after four original members: *MCMI*, *AGAMOUS*, *DEFICIENS*, and *SRF*.

A-Function Genes

In Arabidopsis, the A-class is mainly represented by *APETALA1* (AP1) and *APETALA2* (AP2).

- AP1 is a MADS-box gene found in the outer two whorls, determining sepal and petal identity. It also functions in the transition of floral meristem identity.
- AP2 is distinct from the majority of other floral homeotic genes as it belongs to the AP2/ERF transcription factor family instead of being a MADS-box gene. AP2 suppresses the expression of the C-function gene (*AGAMOUS*) in the outer whorls, preserving the A-domain.
- In *Antirrhinum majus*, the corresponding A-class genes consist of *SQUAMOSA* and *LIPLESS1/2*.

B-Function Genes

The B-function genes act in the second and third whorls, specifying petals and stamens. They include:

- **APETALA3 (AP3)** and **PISTILLATA (PI)** in *Arabidopsis*.

- Their *Antirrhinum* homologs are *DEFICIENS (DEF)* and *GLOBOSA (GLO)*. AP3 and PI proteins form heterodimers, which are essential for their stability and DNA-binding ability. The AP3/PI complex activates downstream genes that lead to petal and stamen differentiation. Mutations in either gene lead to the loss of B-function, causing petals to convert to sepals and stamens to carpels.

C-Function Genes

The C-function is represented by **AGAMOUS (AG)** in *Arabidopsis* and its *Antirrhinum* counterpart *PLENA (PLE)*.

AG is expressed in the inner two whorls (stamens and carpels) and plays a dual role:

1. Specifies stamen and carpel identity.
2. Terminates floral meristem activity, preventing indeterminate flower formation.

Mutations in *AG* result in flowers where stamens are replaced by petals and carpels by a new flower, creating a reiterative floral structure.

Genetic Interactions and Regulation

The ABC genes are regulated both spatially and temporally through complex genetic networks. The expression domains of A, B, and C genes are established by upstream regulators such as **LEAFY (LFY)** and **WUSCHEL (WUS)**, which activate floral meristem identity and C-class expression respectively.

- **LFY** promotes the expression of *API*, *AP3*, *PI*, and *AG*, linking meristem identity to organ identity.
- **WUS** acts in the center of the floral meristem to activate *AG*, but its expression is terminated once *AG* becomes active — a feedback loop that ensures meristem determinacy.

The mutual repression between A and C genes is fundamental:

- A-function genes (like *AP2*) repress *AG* in outer whorls.
- C-function gene (*AG*) represses *API/AP2* in inner whorls. This mutual exclusion creates distinct expression domains that define organ identity.

CONCLUSION

The ABC model of flower development is one of the most important achievements in plant developmental biology, explaining how flowers form their distinct reproductive organs through the coordinated action of homeotic genes. Studies in *Arabidopsis thaliana* and *Antirrhinum majus* demonstrated that combinations of A, B, and C class genes determine the identity of sepals, petals, stamens, and carpels in specific floral whorls. Most of these genes encode MADS-box transcription factors that regulate organ differentiation and floral meristem activity through precise spatial and temporal gene expression. The model also revealed the importance of genetic interactions, particularly the mutual repression between A and C functions, which ensures proper floral patterning. Beyond explaining floral organogenesis, the ABC model laid the foundation for advanced concepts such as the ABCDE and floral quartet models. Its significance extends from evolutionary biology to modern agriculture, where understanding floral genetics supports crop improvement, hybrid seed production, and ornamental plant breeding. Overall, the ABC model remains a landmark framework for understanding the molecular basis of flower development and diversity in angiosperms.

REFERENCES

- Bowman, J. L., & Moyroud, E. (2024). Reflections on the ABC model of flower development. *The Plant Cell*, 36(5), 1334–1357.
- Bowman, J. L., Smyth, D. R., & Meyerowitz, E. M. (2012). The ABC model of flower development: Then and now. *Development*, 139(22), 4095–4098.
- Causier, B., Schwarz-Sommer, Z., & Davies, B. (2010). Floral organ identity: 20 years of ABCs. *Seminars in Cell & Developmental Biology*, 21(1), 73–79.
- Ditta, Gary; Pinyopich, Anusak; Robles, Pedro; Pelaz, Soraya; Yanofsky, Martin F. (2004). "The SEP4 Gene of

- Arabidopsis thaliana* Functions in Floral Organ and Meristem Identity". *Current Biology*. **14** (21): 1935–40.
- Favaro, R; Pinyopich, A; Battaglia, R; Kooiker, M; Borghi, L; Ditta, G; Yanofsky, MF; Kater, MM; Colombo, L (2003). "MADS-box protein complexes control carpel and ovule development in *Arabidopsis*". *The Plant Cell*. **15** (11): 2603–11.
- Irish, V. F. (2017). The ABC model of floral development. *Current Biology*, **27**(17), R887–R890.
- Kanno, Akira; Saeki, Hiroshi; Kameya, Toshiaki; Saedler, Heinz; Theissen, Günter (2003). "Heterotopic expression of class B floral homeotic genes supports a modified ABC model for tulip (*Tulipa gesneriana*)". *Plant Molecular Biology*. **52** (4): 831–41.
- Krogan, Naden (November 2012). "APETALA2 negatively regulates multiple floral organ identity genes in *Arabidopsis* by recruiting the co-repressor TOPLESS and the histone deacetylase HDA19". *Development*. **139** (22): 41 80–90.
- Nakamura, Toru; Fukuda, Tatsuya; Nakano, Masaru; Hasebe, Mitsuyasu; Kameya, Toshiaki; Kanno, Akira (2005). "The modified ABC model explains the development of the petaloid perianth of *Agapanthus praecox* ssp. *Orientalis* (Agapanthaceae) flowers". *Plant Molecular Biology*. **58** (3): 435–45.
- Ó'Maoiléidigh, D. S., Graciet, E., & Wellmer, F. (2014). Gene networks controlling *Arabidopsis thaliana* flower development. *New Phytologist*, **201**(1), 16–30.
- Pelaz, S., Ditta, G. S., Baumann, E., Wisman, E., & Yanofsky, M. F. (2000). B and C floral organ identity functions require SEPALLATA MADS-box genes. *Nature*, **405**(6783), 200–203.
- Prunet, N., & Jack, T. P. (2014). Flower development in *Arabidopsis*: There is more to it than learning your ABCs. *Methods in Molecular Biology*, **1110**, 3–33.
- Rijpkema, A. S., Vandenbussche, M., Koes, R., Heijmans, K., & Gerats, T. (2010). Variations on a theme: Changes in the floral ABCs in angiosperms. *Seminars in Cell & Developmental Biology*, **21**(1), 100–107.
- Soltis, D.E., Chanderbali, A. S., Kim, S., Buzgo, M., & Soltis, P. S. (2010). The ABC model and its applicability to basal angiosperms. *Annals of Botany*, **100**(2), 155–163.
- Sommer, H; Beltrán, JP; Huijser, P; Pape, H; Lönnig, WE; Saedler, H; Schwarz-Sommer, Z (1990). "Deficiens, a homeotic gene involved in the control of flower morphogenesis in *Antirrhinum majus*: The protein shows homology to transcription factors". *The EMBO Journal*. **9** (3): 605–13.
- Vandenbussche, M., Zethof, J., Royaert, S., Weterings, K., & Gerats, T. (2004). The duplicated B-class heterodimer model: whorl-specific effects and complex genetic interactions in *Petunia hybrida* flower development. *The Plant Cell*, **16**(3), 741–754.
- Wellmer, F., Graciet, E., & Riechmann, J. L. (2014). Specification of floral organs in *Arabidopsis*. *Journal of Experimental Botany*, **65**(1), 1–9.