



## From Weather Data to Farm Decisions: The Role of AI in Modern Agrometeorology

**Moinuddin<sup>1</sup>**  
**Prerna Negi<sup>2</sup>**  
**Sarthak Verma<sup>3</sup>**

<sup>1</sup>Professor, Department of  
Agronomy, Shri Guru Ram Rai  
University, Dehradun.

<sup>2,3</sup>Research Scholar, Department  
of Agronomy, Shri Guru Ram  
Rai University, Dehradun.



\*Corresponding Author  
**Moinuddin\***

### Article History

Received : 21.08.2026


Revised : 26.08.2026

Accepted : 31.08.2026

This article is published under the  
terms of the [Creative Commons  
Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/).

### INTRODUCTION

Agriculture has always been weather dependent, but climate change is making that link harder to manage. Rainfall is less predictable, heatwaves are more common, dry spells hit crops in their vulnerable stages, and even a few hours of intense storms can wipe out months of investment. Conventional weather reports remain useful, they are frequently handwaved at the farmer as nothing more than big city advice with no bearing on the farm. AI is enabling tomorrow's agrometeorology, which transforms a complex array of weather-related data into near real-time, geospatial recommendations for farmers. So its real value is not just in producing a forecast, but in linking atmospheric observations to crop, water, pest, labour, machinery and risk management decisions.



### Building Intelligence from Diverse Data

Modern agrometeorology uses information derived from weather stations, satellites, doppler radar, soil probes, crop models, UAVs, farm records and mobile phones. However, none of the sources really give the complete picture for agriculture. Local temperature, humidity, rainfall and wind are measured at weather stations, and cloud movement, vegetation status, surface temperature and soil-moisture patterns over larger areas can be observed by satellites.

Artificial intelligence integrates these diverse data streams, uncovers relationships among them and dynamically updates predictions as new observations are obtained. This integration enables advisory systems to go beyond generic weather forecasts and predict how the expected weather conditions might affect a crop, growth stage, soil type or management practice at a particular location.

### **A New Generation of Weather Forecasting**

Conventional numerical weather prediction employs physical equations to model atmospheric changes. They remain crucial, but require massive amounts of computing and can still possess regional biases. Machine-learning models can infer atmospheric patterns from a history of observations and model analyses spanning decades, making forecasts quickly, and correcting systematic errors. In February 2025, the European Centre for Medium-Range Weather Forecasts (ECMWF) brought its Artificial Intelligence Forecasting System (AIFS) operational alongside its physics-based system. ECMWF reported improvements for a number of forecast metrics including tropical-cyclone tracks, to illustrate that AI forecasting has moved from experimentation to operational meteorology. The best of the future models will integrate physical knowledge, observations, and data-based learning rather than regard them as competing methodologies.

### **From Regional Forecasts to Field-Level Information**

A district-level prediction maybe not represent the farms divided by the heterogeneous landscape. Assisted downscaling via AI applies information on topography, land cover, elevation and historic weather data to produce finer-scale local predictions, and bias correction compensates systematic forecast errors with observed data. With the use of weather stations and field sensors, these techniques enhance the accuracy in the estimation of rainfall, temperature, humidity, wind and evapotranspiration.


However, hyperlocal predictions must be validated rigorously, as the boom in sensors is leading to the introduction of some very poorly maintained sensors.

### **Choosing the Right Time to Sow**

The sowing date is one of the most important decisions in rainfed farming. Sowing immediately after an isolated early shower may result in seeds being subjected to a desiccating spell of weather, whereas late sowing can lead to shortened growing period and lower yield. AI models can take into account rainfall onset, soil moisture, seasonal forecasts, crop duration and the likelihood of consecutive dry days to detect a less risky time for planting. Advisories can be differentiated even at the level of crop, variety and soil type as the needs for establishment vary widely. An intelligent advisory, instead of simply informing that monsoon rainfall has started, should advise on sowing a short-duration variety; waiting for one more such effective rainfall or preparing land for sowing while preserving moisture in the soil.

### **Smarter and More Efficient Irrigation**

Scheduling irrigation based on weather information has traditionally been a function of the reference evapotranspiration and crop coefficients. AI can further enhance this method by considering forecast temperature, radiation, humidity, wind, probability of rainfall, soil-moisture data and crop-development information. The based system is able to predict the timing and quantity of irrigation as well as whether the irrigation operation should be postponed due to an expected rain. It could also detect spatial variability in the field and facilitate variable irrigation if appropriate machinery is on hand. These functions are particularly critical for the dry areas of the world where over irrigation can waste limited water and energy supplies, and where under irrigation of flowering, fruiting or grain filling can cause irreversible yield and quality losses.



### Weather-Informed Pest and Disease Management

Temperature, humidity, rainfall, wind and leaf wetness have the strongest effect on insects, pathogens and their host plants. AI models could integrate these variables with pest-surveillance data, crop stage, prior outbreaks and satellite data to provide predictions of insect or disease risk. A disease warning may be able to predict the existence of conditions favorable for the germination of spores prior to widespread dissemination of visible symptoms, insect models can forecast emergence, migration or explosive population growth temperature-based phenology. This advice enables farmers to examine fields at the optimal time and to treat only if risk warrants taking action. Crucially, weather-based alerts should prompt field scouting, not automatic pesticide use, as current pest populations, natural enemies and the economic threshold must continue to be part of the decision.

### Preparing for Extreme Weather

Extreme weather warnings are useful only if they are associated with actions people can

take to protect themselves. AI can be used to convert predictions of heat, frost, hail, heavy rain, high winds or drought into estimates of crop damage and financial exposure. A heat alert could lead to additional irrigation, shade manipulation or altered harvesting timing, while a freeze warning might justify protective irrigation, orchard heating or postponed transplanting. Predictions of heavy rain, for example, help farmers prepare their drains, delay application of fertilizer and move machinery, livestock or harvested crops to higher ground. More and more research promotes multi-hazard early-warning systems that integrate meteorological and geospatial information, that communicate uncertainty and that incorporate feedback from the people who should act on the warning.

### Supporting Yield and Harvest Decisions

Artificial intelligence can link weather history and crop conditions with predicted yield, maturity and harvest date. Temperature, rainfall, radiation, soil moisture, vegetation indices, and management information may be integrated in models for monitoring crop growth during the season. With the updated

estimates of yield, the farmers get a better opportunity to plan their labour, machinery, storage, transportation and marketing etc. Weather-sensitive harvest guidance holds particular importance for fruits, vegetables, grains and seed crops, as rainfall or very high humidity can reduce quality and amplify postharvest loss. Yet yield forecasts are in a sense expected to be refined estimates rather than definitive solutions, as unexpected pest, nutrient restrains, localized storms or management differences can change end production even if seasonal weather is perfectly represented.

### **Delivering Advice Farmers Can Use**

Scientific accuracy does not necessarily lead to better agricultural decisions. Advice based on forecasts needs to be timely, understandable and appropriate to the resources of farmers. AI platforms can transform forecasts into personalized messages that reach farmers via apps, SMS, voice services, chatbots, radio and extension networks. Instructions can be tailored by location, crop, variety, growth stage, irrigation and language. Actionable advisories include clear information on the forecast time period, expected severity, recommended action and remaining uncertainty.

### **Opportunities for Indian Agriculture**

The highly diverse farm conditions in India also call for agrometeorological services from a location-specific perspective. The impact of a single weather event on irrigated wheat, rainfed pulses, protected vegetables and fruit orchards can also vary. AI can help integrate forecasts with local crop calendars, soil conditions, and management options to enhance advisories on sowing, irrigation, spraying, harvesting and livestock protection. Recent climate information led projects in Odisha show how local production risk management services can be upscaled to reach thousands of farmers. To scale up such systems, stronger networks of automatic-

weather-stations and reliable village-level observations will be required, as well as cooperation between meteorological agencies, agricultural universities and extension departments, and ongoing feedback from farmers as to whether the recommendations were understandable, timely and practically useful.

### **Trust, Explainability and Uncertainty**

Agricultural action should never be taken on a prediction without explanation that is given as a certainty. Weather and crop systems already have a level of inherent unpredictability, and AI models may break down if they are faced with data far outside their training set. Probabilities, alternative scenarios, and confidence levels should be available from reliable platforms, too. Explainable AI can tell which variables were the largest contributors to a warning, providing meteorologists, extension personnel, and farmers a means to evaluate if the outcome makes sense. Latest science demonstrates that expert-informed AI can probabilistically detect agricultural risks and simultaneously reveal their principal climatic drivers. Before a model is trusted with high-stakes farm decisions, it must be independently validated, compared to traditional approaches, and transparently report its errors.

### **Inclusion, Ownership and Responsible Use**

The advantages of AI will continue to be skewed if the delivery of digital services relies on costly sensors, constant internet connectivity, or data that can only be obtained from aggregators of large commercial farms. Rural connectivity, open meteorological datasets, local-language services and technical capacity of extension staff require public investment. Farmers should be informed what data is being collected, for what purpose, who has access to it, and if that data can be used commercially. Privacy should be protected in responsible systems, farmer data rights should be acknowledged, and recommendations

should not be biased by commercial interests that are not disclosed. Participatory design is equally crucial: farmers, women, tenant farmers, local specialists and other disenfranchised groups should participate in deciding which risks are tracked, and on which advisories formats are truly effective.

### **The Continuing Role of Human Expertise**

AI does not replace the need for agrometeorologists, crop experts, or extension agents. Instead, it redefines their role from undertaking the manual processing of atomised data to the interpretation of model outcomes, the assessment of their relevance to the actual farming situation and the support to farm-level decision making in the presence of uncertainty. Local experts can identify failed sensors, bizarre crop responses, and cultural limitations that a mechanistic model can't account for. Farmers, too, have intimate understanding of soils and microclimates as well as traditional indicators of risk and risk tolerance that are scarcely captured in official datasets. The most reliable advice system is thus a continuous learning system: observations feed back into models, models

produce guidance, experts assess the guidance, farmers act or adapt, and their experiences feedback to improve future recommendations. Human control makes automated prediction an accountable service to agriculture.

### **CONCLUSION**

The future of agrometeorology is in transitioning from weather observing to decision intelligence. Artificial intelligence has the potential to combine vast amounts of heterogeneous data, increase the speed of forecasting, yield field-specific risk estimates, and provide personalized guidance at the point of decision making. It will not be judged on algorithmic complexity but on better planting decisions, efficient irrigation, timely pest surveillance, less weather damage and more stable farm income. These results depend on good observations, transparent models, reliable communication, strong extension and responsible data governance. Once these are in place, AI is not simply a forecasting tool; it is a practical link between atmospheric science and resilient, resource-efficient and farmer-centred agriculture.