## ISSN (E): 2583 - 1933

Available online at http://currentagriculturetrends.vitalbiotech.org/

Curr. Agri.Tren.: e- Newsletter, (2024) 3(10), 35-37

Article ID: 346



# **Cross-Talk Between Plant Hormones in Disease Response**

# Sonakshi<sup>1</sup>, Jatin Badwal<sup>2</sup> and Ishwar Singh<sup>2</sup>

<sup>1</sup>Department of Plant Pathology, CCS Haryana Agricultural University, Hisar- 125004 <sup>2</sup>Department of Agriculture, Maharishi Markandeshwar (Deemed to be University) Mullana, Ambala-133207, Haryana



**Article History** Received: 13. 10.2024 Revised: 16. 10.2024 Accepted: 20. 10.2024

This article is published under the terms of the <u>Creative Commons</u> <u>Attribution License 4.0</u>.

### INTRODUCTION

Plants, as sessile organisms, are constantly exposed to diverse biotic stresses such as pathogens, pests, and parasitic organisms. To survive, they have developed sophisticated defense mechanisms mediated by a complex network of signaling pathways. Central to this defense system is the interplay or "cross-talk" between plant hormones. Key hormones, including salicylic acid (SA), jasmonic acid (JA), ethylene (ET), abscisic acid (ABA), and auxins, work in an interconnected manner to fine-tune immune responses. Understanding the crosstalk between these hormonal pathways is vital for advancing agricultural practices and developing disease-resistant crops.

#### Key Plant Hormones in Disease Response

#### • Salicylic Acid (SA)

SA is crucial for systemic acquired resistance (SAR), a long-lasting immune response triggered by localized pathogen infection. It is primarily effective against biotrophic pathogens, which derive nutrients from living host cells (Klessig *et al.*, 2018).

#### • Jasmonic Acid (JA)

JA, along with its derivatives like methyl jasmonate (MeJA), mediates defenses against necrotrophic pathogens and herbivorous insects. It triggers the expression of defense-related genes and promotes secondary metabolite production (Wasternack & Song, 2017).

#### • Ethylene (ET)

ET is often involved in modulating JA signaling and inducing defenses against necrotrophic pathogens. It also plays a role in the regulation of cell death during hypersensitive responses (Broekgaarden *et al.*, 2015).

#### • Abscisic Acid (ABA)

Traditionally recognized for its role in abiotic stress tolerance, ABA also modulates plant-pathogen interactions. Its role in defense is complex, sometimes promoting susceptibility by antagonizing SA or JA pathways (Cao *et al.*, 2011).



Available online at http://currentagriculturetrends.vitalbiotech.org

#### Auxins, Cytokinins, and Gibberellins

These growth hormones can influence disease resistance indirectly. For example, auxin signaling can promote susceptibility to certain pathogens by repressing immune signaling (Kazan & Manners, 2009).

#### **Mechanisms of Hormonal Cross-Talk**

#### • SA and JA: Antagonistic Interactions

SA and JA are well-documented for their antagonistic relationship. SA signaling, effective against biotrophic pathogens, often suppresses JA signaling, which is more effective against necrotrophs. This antagonism is mediated by transcriptional regulators such as NPR1 (Non-expressor of Pathogenesis-Related Genes 1) (Spoel *et al.*, 2007). For example, Arabidopsis infected with *Pseudomonas syringae* exhibits suppressed JA responses due to SA-mediated repression.

#### • JA and ET: Synergistic Interactions

JA and ET often work synergistically to combat necrotrophic pathogens. The coregulation of defense genes like *PDF1.2* in Arabidopsis is a hallmark of this interaction. ET amplifies JA signals, enhancing the plant's ability to mount a robust defense against pathogens like *Botrytis cinerea* (Broekgaarden *et al.*, 2015).

#### • ABA and SA/JA: Complex Interplay

ABA can both suppress and enhance immune responses, depending on the context. It can antagonize SA signaling, making plants more susceptible to biotrophic pathogens. Conversely, it may synergize with JA to enhance defenses against necrotrophs (Cao *et al.*, 2011). This duality highlights the need for spatiotemporal regulation of ABA signaling during pathogen attacks.

# • Emerging Roles of Cytokinins and Auxins

Cytokinins have been shown to modulate immune responses by balancing SA and JA signaling. Conversely, auxins are often exploited by pathogens to suppress immune signaling, promoting disease (Kazan & Manners, 2009). **Molecular Players in Hormonal Cross-Talk** Several key proteins and transcription factors mediate hormonal interactions:

- NPR1: Central in SA-JA antagonism, integrating signals to fine-tune defenses.
- ERF (Ethylene Response Factors): Mediate JA-ET synergistic responses.
- WRKY Transcription Factors: Regulate the expression of defense genes in response to hormonal cues (Broekgaarden *et al.*, 2015).

#### **Applications and Future Perspectives**

The knowledge of hormonal cross-talk can be harnessed to develop disease-resistant crops. For example:

- **Biotechnology**: Engineering plants to modulate specific hormone pathways can enhance resistance to multiple pathogens. For instance, boosting SA signaling could improve resistance against biotrophs.
- Sustainable Agriculture: Hormonal priming using natural compounds like JA derivatives can prepare plants for impending pathogen attacks without genetic modifications (Wasternack & Song, 2017).

#### Future research should focus on:

- 1. Identifying novel molecular players in hormonal interactions.
- 2. Understanding the role of secondary messengers, such as reactive oxygen species (ROS), in hormone-mediated defense.
- 3. Exploring how environmental factors influence hormonal cross-talk.

#### CONCLUSION

The cross-talk between plant hormones forms the backbone of plant immunity, enabling plants to prioritize and tailor their defenses against diverse pathogens. While the



Available online at http://currentagriculturetrends.vitalbiotech.org

antagonistic and synergistic interactions between hormones like SA, JA, and ET are well-studied, ongoing research continues to unravel the complexity of these signaling networks. Leveraging this knowledge in agriculture could revolutionize crop protection strategies, ensuring food security in the face of escalating biotic stresses.

#### REFERENCES

- Broekgaarden, C., Caarls, L., Vos, I. A., Pieterse, C. M., and Van Wees, S. C. (2015). Ethylene: Traffic controller on hormonal crossroads to defense. *Plant Physiology*, 169(1), 237-246.
- Cao, F. Y., Yoshioka, K., and Desveaux, D. (2011). The roles of ABA in plantpathogen interactions. *Journal of Plant Research*, 124(4), 489-499. https://doi.org/10.1007/s10265-011-0402.

- Kazan, K., & Manners, J. M. (2009). Linking development to defense: Auxin in plant-pathogen interactions. *Trends in Plant Science*, 14(7), 373-382.
- Klessig, D. F., Tian, M., and Choi, H. W. (2018). Multiple targets of salicylic acid and its derivatives in plants and animals. *Frontiers in Immunology*, 9, 206.
- Spoel, S. H., Johnson, J. S., and Dong, X. (2007). Regulation of tradeoffs between plant defenses against pathogens with different lifestyles. *Proceedings of the National Academy* of Sciences, 104(47), 18842-18847.
- Wasternack, C., and Song, S. (2017). Jasmonates: Biosynthesis, metabolism, and signaling by proteins activating and repressing transcriptions. *Journal of Experimental Botany*, 68(6), 1303-1321.