Available online at
http://currentagriculturetrends.vitalbiotech.org/

ISSN (E): 2583 – 1933

Curr. Agri.Tren.:e- Newsletter, (2026) *5*(5), 82-88

Current Agriculture Trends: e-Newsletter

Article ID: 641

Germplasm Conservation in Fruit Crops: Integrating Traditional Knowledge and Modern Science for Food Security

Meena Kumari[1], Savita Kumari[2], Rimpika[3], Kiran Thakur[4] and Shilpa[5]

[1]Assistant Professor, College of Horticulture and Forestry, YSPUH&F, Nauni, Solan, Himachal Pradesh (173 230), India
[2]Subject Matter Specialist, College of Horticulture and Forestry, Neri, Hamirpur, Himachal Pradesh (177 001), India
[3]Assistant Professor, Litchi and Mango Research Station, Nagrota Bagwan, Kangra, Himachal Pradesh (176 047), India
[4]Assistant Professor, Directorate of Extension, College of Horticulture and Forestry, YSPUH&F, Nauni, Solan, Himachal Pradesh (173 230), India
[5] Assistant Professor, Department of Fruit Science, Regional Horticultural Research & Training Station, Dhaulakuan

Open Access

*Corresponding Author
Meena Kumari*

Article History
Received: 5. 5.2026
Revised: 10. 5.2026
Accepted: 15. 5.2026

This article is published under the terms of the Creative Commons Attribution License 4.0.

INTRODUCTION

Germplasm conservation in fruit crops is fundamental to sustainable agriculture and plays a vital role in ensuring long-term food security, nutritional diversity, and ecological stability in the face of rapid environmental and socio-economic changes. Fruit crops are important sources of nutrition, income, and cultural value, and their continued productivity depends on the availability of a wide genetic base capable of adapting to changing conditions. However, this genetic diversity is increasingly threatened by climate change, intensive agricultural practices, emerging pests and diseases, habitat loss, urbanization, and the growing replacement of traditional varieties with genetically uniform, high-yielding cultivars. Such factors contribute to the erosion of valuable genetic resources and weaken the resilience of agricultural systems against future challenges. In this scenario, the conservation of fruit crop germplasm has become critically important. Germplasm refers to the total genetic material present within a species, including wild relatives, landraces, traditional cultivars, and improved varieties. These genetic resources possess valuable traits such as drought tolerance, disease resistance, enhanced nutritional quality, and adaptability to adverse environmental conditions. Conserving this diversity requires an integrated approach that combines traditional knowledge with modern scientific technologies. Indigenous practices such as farmer-led selection, seed exchange systems, and community-based conservation have long contributed to the maintenance of on-farm diversity and continue to play a significant role in preserving genetic resources. These traditional methods are now supported by advanced technologies such as cryopreservation for long-term storage of genetic material, molecular characterization for assessing genetic variability, and genomic tools that enhance the effective utilization of germplasm in crop improvement programs.

Both in situ and ex situ conservation strategies are essential and complementary for the effective preservation of fruit crop diversity. In situ conservation involves maintaining plant genetic resources within their natural habitats and farming systems, thereby allowing continuous evolution and adaptation. This includes the conservation of wild fruit species in natural ecosystems as well as on-farm conservation practiced by farmers. Ex situ conservation, in contrast, involves preserving genetic material outside its natural environment through methods such as field gene banks, seed banks, botanical gardens, tissue culture repositories, and in vitro conservation systems. Although each method has specific advantages and limitations, their combined application provides a comprehensive framework for safeguarding valuable genetic resources for present and future generations. Despite advancements, germplasm conservation faces numerous challenges, including high costs of maintenance, technical limitations, lack of standardized protocols, insufficient documentation, and issues related to access, ownership, and benefit-sharing of genetic resources. Addressing these challenges requires strong international collaboration, effective policy frameworks, and adequate funding mechanisms. Global initiatives and partnerships among governments, research institutions, non-governmental organizations, and local communities are crucial to ensure the equitable conservation and utilization of germplasm. Public awareness and education also play a key role in promoting the value of biodiversity and encouraging community participation in conservation efforts. Ultimately, the future of fruit crop germplasm conservation lies in adopting a holistic, multidisciplinary approach that integrates ecological principles, cutting-edge technology, and socio-cultural dimensions. By combining community knowledge with scientific innovation and fostering global cooperation, it is possible to build resilient agricultural systems capable of meeting the food and nutritional needs of a growing population while preserving the rich genetic heritage of fruit crops for future generations.

WILD SPECIES AND DIVERSITY

The role of wild species in the fruit improvement programme is increasingly becoming important as the donor source for many of the disease and pest resistance. However, in most of the perennial trees, wild species or the indigenous germplasm has not been evaluated extensively either by morphological or by molecular means. Some geographical areas may be richer in biodiversity than other areas, and some species may also have more variation than others in a particular area. Conservation of germplasm is very important, because many species are becoming extinct and many others are threatened and endangered. The diversity of some fruits is well documented, while for others relatively little work has been done (Arora, 1994). Gaps in collections are found both between species and between regions. This is especially true for both underutilized species and wild crop relatives, where big gaps are noted. Kostermans and Bompard (1993) indicate that *Mangifera blommesteinii*, *M. leschenaultii*, *M. superb* and. *M. paludosa* are in real danger of extinction. High genetic erosion has been noticed for jackfruit, Citrus spp. and Litchi chinensis in a survey carried out by the International Centre for Underutilized Crops (ICUC) and IPGRI (Haq, 1994). It is to be mentioned here that collection and utilization of wild species is not an easy task, as they require specific climate and do not so easily get acclimatized to the ex-situ conditions on introduction.

Importance of Germplasm Conservation

Germplasm conservation in fruit crops plays a vital role in ensuring sustainable agriculture, food security, and biodiversity preservation. Fruit crops are essential not only for human nutrition but also for the economic stability of many regions around the world. The genetic diversity present in these crops contains valuable traits such as superior taste, enhanced

nutritional value, resistance to pests and diseases, and adaptability to varying environmental conditions. Conserving germplasm helps protect this genetic wealth and ensures that important traits remain available for future crop improvement programs. With increasing challenges such as climate change, emerging diseases, and habitat loss, the conservation of fruit crop germplasm has become more important than ever. It provides breeders and researchers with the resources needed to develop improved varieties capable of tolerating environmental stresses and maintaining productivity under changing conditions. In addition, genetic diversity supports continuous agricultural innovation by enabling the development of new cultivars with better yield, quality, and resilience. Beyond agricultural benefits, germplasm conservation also contributes to the protection of biodiversity, as fruit crops often support various ecosystems by providing food and habitat for many organisms. Therefore, conserving germplasm in fruit crops is essential for maintaining ecological balance, strengthening agricultural sustainability, and securing a diverse and nutritious food supply for present and future generations.

Need for germplasm conservation of endangered plants

There are several reasons why breeders use this technique which is as follows;

- Genetic diversity among plant species is rapidly declining due to environmental and human-related factors.
- Humans and animals depend heavily on plants for food, as major crops such as wheat, maize, and rice form the basis of their nutrition.
- Plants are also essential for various human activities, including construction materials, production of fibers, waxes, perfumes, medicines, and other useful products.
- Deforestation and habitat destruction have endangered many valuable plant varieties, creating an urgent need for their conservation and preservation.
- Genetic diversity helps maintain ecosystem stability by supporting sustainable food chains and ecological balance, which can be preserved through germplasm conservation.
- conserving plant biodiversity also enhances the aesthetic and ecological value of natural ecosystems, contributing to environmental sustainability.

IN-SITU CONSERVATION:

In-situ conservation is on site conservation or the conservation of genetic resources in natural populations of plant or animal species, such as forest genetic resources in natural populations of tree species. It is the process of protecting an endangered plant or animal species in its natural habitat, either by protecting or cleaning up the habitat itself, or by defending the species from predators. It is applied to conservation of agricultural biodiversity in agro forestry by farmers, especially those using unconventional farming practices. In-situ conservation is being done by declaring area as protected area.

In India following types of natural habitats are being maintained:

1. National Park
2. Wildlife sanctuaries
3. Biosphere reserve

National Park

A national park is an area which is strictly reserved for the betterment of the wildlife and where activities like forestry, grazing on cultivation are not permitted. In these parks, even private ownership rights are not allowed. Their boundaries are well marked and circumscribed. They are usually small reserves spreading in an area of 100.

WILDLIFE SANCTUARIES:

A sanctuary is a protected area which is reserved for the conservation of only animals and human activities like harvesting of timber, collecting minor forest products and private ownership rights are allowed as long as they do not interfere with well-being of animals. Boundaries of sanctuaries are not well defined

and controlled biotic interference is permitted, e.g., tourist activity.

BIOSPHERE RESERVES:

It is a special category of protected areas where human population also forms a part of the system. They are large protected area of usually more than 5000 sq.km. A biosphere reserves has 3 parts- core, buffer and transition zone.

1. Core zone is the inner zone; this is undisturbed and legally protected area.
2. Buffer zone lies between the core and transition zone. Some research and educational activities are permitted here
3. Transition zone is the outermost part of biosphere reserves. Here cropping, forestry, recreation, fishery and other activities are allowed.

EX-SITU CONSERVATION

Ex situ conservation of fruit crops involves preserving their genetic diversity outside their natural habitats. This method typically includes storing seeds, tissues, or living collections in controlled environments such as gene banks, botanical gardens, or orchards. These repositories serve as invaluable reservoirs, protecting diverse fruit varieties from potential threats, including climate change, diseases, or habitat loss. Ex situ conservation facilitates research, breeding programs, and public awareness by providing accessible collections of fruit crops .Orchestrating a strategic blend of seed banks, living collections, and advanced technologies, ex situ conservation acts as a vital insurance policy, ensuring the continued existence and utilization of diverse and resilient fruit crop genetic resources.

***IN-VITRO* CONSERVATION**

Two basic approaches are followed to maintain germplasm collections in-vitro:

(i) Minimal growth, and (ii) Cryopreservation **(Scowcroft, 1984).** Minimal growth conditions for short to medium term storage can be followed in several ways – reduced temperature and/or light; incorporation of sub-lethal levels of growth retardants, induction of osmotic stress with sucrose or mannitol, and maintenance of cultures at a reduced nutritional status particularly reduced carbon, reduction of gas pressure over the cultures, desiccation and mineral oil overlay. The advantage of this approach is that cultures can be readily brought back to normal culture conditions to produce plants on demand. However, the need for frequent sub culturing may pose a great disadvantage including contamination of cultures as well as imposition of selection pressure with subsequent change in genetic make-up due to soma clonal variation. Cryopreservation at the temperature of liquid nitrogen (LN) (-196° C) offers the possibility for long-term storage with maximal phenotypic and genotypic stability (Steponkus, 1985). This method being relatively convenient and economical, large number of genotypes and variants could be conserved and thus maximize the potential for storage of genetically desirable material.

Seed banks: Ex situ conservation through seed banks is a pivotal strategy in safeguarding the genetic diversity of fruit crops. Seed banks, as repositories for storing seeds under controlled conditions, play a critical role in preserving the genetic material of various plant species, including fruits. In the context of fruit crops, seeds encapsulate the genetic information necessary for the growth and development of the next generation, representing a compact and resilient form of genetic storage. Seed banks carefully curate collections of fruit crop seeds, maintaining a diverse array of varieties that might be endangered or rare. By storing seeds at low temperatures, often in sub-zero conditions, seed banks slow down metabolic processes, extending the longevity of the seeds and ensuring the preservation of their genetic integrity. This method allows for long-term storage, providing a backup against unforeseen events such as natural disasters, diseases, or climate-related challenges. Furthermore, seed banks often serve as repositories for wild relatives of fruit crops, enriching the genetic diversity available for future breeding

programs. The systematic organization and cataloging of seeds in seed banks facilitate research and the development of new varieties, contributing to the resilience and adaptability of fruit crops in the face of evolving agricultural challenges. Ex situ conservation via seed banks thus acts as a biological insurance policy, safeguarding the genetic resources that underpin the future sustainability and productivity of fruit crops.

Field gene banks: Ex situ conservation through field gene banks emerges as a dynamic and living strategy for preserving the genetic diversity of fruit crops. Unlike traditional seed banks, field gene banks maintain living collections of fruit crops, often organized in orchards, botanical gardens, or designated agricultural fields. These gene banks house a wide array of fruit varieties, including both cultivated and wild relatives, allowing for the maintenance of genetic diversity in an environment that mirrors real-world conditions. By cultivating fruit crops in these controlled settings, field gene banks provide a more dynamic approach to germplasm conservation, allowing for the observation of plant characteristics, growth patterns, and interactions with the environment. This living collection serves as a valuable resource for researchers, breeders, and farmers seeking to understand and utilize the genetic potential of diverse fruit varieties. Field gene banks also play a role in public awareness and education, providing a tangible and accessible display of the richness of fruit crop biodiversity. Through the cultivation and study of these living collections, field gene banks contribute to the identification of traits crucial for enhancing the resilience, adaptability, and sustainability of fruit crops in changing agricultural landscapes. In this way, ex situ conservation via field gene banks bridges the gap between in situ preservation and traditional seed banks, offering a holistic and interactive approach to maintaining the genetic resources essential for the future of fruit crop agriculture.

Cryopreservation: Ex situ conservation through cryopreservation represents a cutting-edge and highly effective method for safeguarding the genetic diversity of fruit crops. Cryopreservation involves preserving plant genetic material, such as seeds, embryos, or tissues, at extremely low temperatures, usually in liquid nitrogen. In the context of fruit crops, this technique offers several advantages. By arresting metabolic activity and slowing cellular processes to a near halt, cryopreservation allows for the long-term storage of genetic material without the risk of deterioration. This is particularly integrity of seeds and tissues of fruit crops over extended periods. Cryopreservation also minimizes the risk of genetic changes or mutations, ensuring that the stored germplasm remains true to its original genetic composition. The method is especially valuable for conserving recalcitrant seeds, which are susceptible to damage during traditional storage methods. Moreover, cryopreservation facilitates the storage of a diverse range of fruit crop varieties, including those that are rare, endangered, or difficult to maintain through other conservation methods. It has saved over ten thousand species. As an advanced and versatile technique, cryopreservation stands at the forefront of ex situ conservation efforts, offering a sophisticated means of preserving the genetic resources critical for the continued resilience and adaptability of fruit crops in the face of evolving environmental challenges.

Tissue culture: Ex situ conservation through tissue culture stands as a sophisticated and precise method for preserving the genetic diversity of fruit crops. Tissue culture involves the *in vitro* cultivation of plant cells, tissues, or organs under controlled conditions. In the context of fruit crops, this technique offers unique advantages for germplasm conservation. By carefully selecting and culturing specific plant tissues, such as shoot tips, embryos, or meristem, tissue culture enables the regeneration of entire plants while maintaining their genetic integrity. This method is particularly beneficial for fruit crops

with recalcitrant seeds or those prone to genetic variations during traditional seed storage. Tissue culture also allows for the rapid multiplication of disease-free and genetically uniform plant material, ensuring the efficient propagation of valuable and often rare fruit varieties. The controlled environment of tissue culture facilities minimizes the risk of contamination and provides optimal conditions for the growth of plant cells .This precision in maintaining genetic characteristics is crucial for preserving the unique traits, such as flavor profiles, nutritional content, and disease resistance that are inherent in diverse fruit crop varieties.

FUTURE PERSPECTIVES

The future of germplasm conservation in fruit crops holds great promise as researchers, scientists, and conservationists navigate the complexities of a rapidly changing agricultural landscape. One key aspect of future perspectives is the integration of traditional knowledge with cutting-edge technologies. Incorporating the wisdom of local communities, often the stewards of diverse fruit varieties, enhances the effectiveness and sustainability of conservation efforts. Additionally, advancements in molecular biology, genomics, and biotechnology offer unprecedented tools for characterizing and utilizing genetic resources. The utilization of high-throughput sequencing, gene editing technologies, and bioinformatics can revolutionize the precision and efficiency of germplasm conservation in fruit crops. Educational initiatives and public awareness campaigns will play a pivotal role in shaping the future of germplasm conservation. By fostering an understanding of the importance of genetic diversity and its impact on food security, policymakers, farmers, and the general public can become active participants in conservation efforts. Furthermore, the emphasis on participatory approaches, where local communities are engaged in decision-making processes, ensures that conservation strategies align with the needs and values of diverse stakeholders. Looking ahead, collaborative international efforts will be essential. Sharing germplasm resources, knowledge, and technologies on a global scale can enhance the resilience of fruit crops against emerging threats and contribute to the development of more robust and adaptable varieties. The future of germplasm conservation in fruit crops lies in a holistic, inclusive, and technologically informed approach, which not only preserves genetic diversity but also paves the way for sustainable and resilient fruit crop agriculture in the face of evolving challenges.

CONCLUSION

In conclusion, germplasm conservation in fruit crops emerges as an indispensable endeavor for ensuring the sustainability, adaptability, and resilience of global agriculture. The significance of preserving the genetic diversity within fruit crops lies in its role as the foundation for breeding programs, offering a reservoir of traits essential for crop improvement. Despite the challenges, from financial constraints to ethical considerations, the imperative to conserve germplasm in fruit crops is underscored by the increasing threats of climate change, emerging diseases, and habitat loss. In navigating these challenges, a multifaceted approach that integrates in situ and ex situ conservation methods, engages local communities, and leverages advanced technologies is paramount. The future of germplasm conservation in fruit crops holds promise through the convergence of traditional knowledge and cutting-edge technologies. Educational initiatives and global collaboration will be pivotal in enhancing public awareness and sharing resources for the sustained preservation of genetic diversity. As we chart the course forward, it is clear that germplasm conservation is not merely a scientific endeavor but a shared responsibility that shapes the future of agriculture, food security, and the rich tapestry of fruit crop biodiversity for generations to come.

Copyright © May, 2026; Current Agriculture Trends

REFERENCES

Haq, N. (1994). *Identification of priority species of tropical fruits for research and development*. Final Report, IPGRI Project No. 93/11 (AOG-APO), International Centre for Underutilized Crops (ICUC) in collaboration with IPGRI-APO, Southampton University, UK.

R. K. Arora (1994). *Plant genetic resources activities at IPGRI office for South Asia*. In *Plant Genetic Resources: National Bureau of Plant Genetic Resources* (pp. 20–22). New Delhi, India.

Scowcroft, W. R. (1984). Genetic variability in tissue culture: Impact on germplasm conservation and plant breeding. In *Plant Biotechnology and Genetic Engineering* Elsevier. (pp. 217–245).

Steponkus, P. L., & Gordon-Kamm, W. J. (1985). Cryoinjury in isolated protoplasts and its relationship to dehydration injury and freezing injury. *Cryo-Letters*, 6, 217–226.