



Botanical Insecticides: Harnessing Plant-Based Solutions for Pest Management

Ishfaq Majeed¹, Ramnivas²
Sandhya M³
Dibyani Nanda⁴
Ningthoujam Samarendra
Singh⁵

¹Ph.D Scholar, Division of
Entomology, SKUAST-K

²Assistant Professor, Department of
Entomology, MSAC, Rahimpur
Uchain, Bharatpur (Raj.)

³Ph.D Scholar, Department of
Entomology, College of Agriculture
UAS, Raichur

⁴M.Sc. Scholar (Ag.), Department of
Entomology, Rani Lakshmi Bai
Central Agricultural University

⁵Scientist, Division of Post-Harvest
Management, ICAR-Central
Institute for Sub-Tropical
Horticulture (ICAR-CISH)
Lucknow, India.



Open Access

*Corresponding Author
Ishfaq Majeed*

Article History

Received: 16.09.2026

Revised: 21.09.2026

Accepted: 26.09.2026

This article is published under the
terms of the [Creative Commons
Attribution License 4.0.](https://creativecommons.org/licenses/by/4.0/)

INTRODUCTION

A crop may be healthy in the morning and then a few days later the insects are multiplying and it is visibly damaged when conditions are conducive. Farmers require the ability to act quickly, but consistent application of the same insecticides may select for resistant pests and may disrupt beneficial organisms. Botanical insecticides provide yet another source of pest control: chemicals plants produce to protect themselves. Neem, pyrethrum and aromatic plants are well-known examples, but new candidates and improved ways to formulate them are being discovered by research. They are useful because they provide growers with more options, even if all of those options are constrained by a tightly integrated pest management program.



How Plants Defend Themselves

Plants cannot run away from insect attack, so many plants produce chemicals that make their leaves, seeds or stems hard to eat. Some of these chemicals may directly repel a pest, or prevent it from laying eggs, or cause it to eat less, or prevent it from growing and reproducing. Some of these chemicals act quickly on the insect nervous system, while others have slower effects that are not visible for several days. A botanical insecticide may be one isolated active ingredient or a mixture of compounds that occur naturally. The differences help to explain the inconsistency among plant-based products in terms of the pests they control and how quickly they work.

How Plants Defend Themselves



Neem: A Widely Studied Example

Neem (*Azadirachta indica*) is one of the better-known sources of botanical insecticides. Azadirachtin, the main insect-active compound of neem, can reduce feeding and interfere with normal development, especially in immature insects. A treated caterpillar may consume less foliage or fail to complete moulting even when it does not die immediately. Neem products have been tested against vegetable, fruit, and field crop pests, but their efficacy is influenced by formulation, application timing, and pest species. Thus, growers should measure success by reduced crop damage and pest development, not by the immediate disappearance of every insect they have exposed to neem.

Pyrethrum and Essential Oils

Pyrethrum derived from the flowers of certain chrysanthemums is appreciated for its fast-acting on several insects. Essential oils of aromatic plants have other potentials including repellency, antifeeding and toxicity to the insects. Oils derived from thymol, eugenol or citronellal-containing substances have received scientific attention, but being effective in laboratory does not mean effective in field. The word “botanical” encompasses a wide range of products that have very different chemical and biological profiles. To choose among them,

one needs to know the pest, the crop, and the purpose, i.e., immediate suppression or prevention of further infestation.

Matching the Product to the Pest

Efficacy of the same treatment is rarely equally good on aphids, whiteflies, caterpillars and stored-grain beetles. How feeding behaviour, protective covering and life stage influence the amount of active ingredient reaching them. For example, a spray targeting insects feeding on the undersides of leaves must have good coverage of those surfaces, and a treatment given after a caterpillar has entered a fruit may be too late. Young larvae are often easier to manage than established populations. Accurate identification and frequent monitoring are critical: they enable growers to select an appropriate product and apply it while the pest is still exposed and crop damage is minimal.

The Challenge of Consistent Performance

The chemical composition of a plant varies with the variety, growing conditions, stage of harvest and the part used. Extraction and storage can introduce additional differences. Therefore, two products with the same plant name may have different concentrations of active material and different effects. Sunlight, heat and rain may further diminish the residual

effect of certain botanical sprays following application. These are among the reasons why good results in the laboratory can sometimes fail to reliably predict control in the field. A commercial product needs standardized ingredients, quality control, appropriate packaging and testing over seasons and locations, not just a plant name.

Better Formulations

Recent research has focused on protecting unstable plant compounds and improving their delivery to pests. Emulsions may facilitate

dispersion of oils in water; encapsulation may shield ingredients and modulate their release. Certain experimental nano emulsions exhibited enhanced dispersion or activity in particular in vitro assays. Although these advances are encouraging, a smaller droplet size alone does not ensure that a product will be effective or safe. Scientists need to evaluate new formulations against conventional ones at realistic rates of application, and on-plant longevity, potential crop damage, cost and impact on non-target organisms. Field validation remains a decisive step.



A Place in Vegetable and Fruit Production

Botanical insecticides may be especially effective on high-value crops where growers need flexible tools for pest management. When applied at the right time, a treatment can help to preserve tender vegetable growth or developing fruit until other control methods kick in. Their typically short persistence may be an advantage in circumstances where long-term residues are not wanted, but shorter persistence may require closer monitoring or a second application. Performance of the product needs to be verified for the specific crop and pest. In cases where trials have measured marketable yield, crop

quality and costs in addition to insect numbers, strong use cases begin to appear, providing growers with the evidence to inform the choices they must make.

Protecting Beneficial Insects

Predators, parasitoids and pollinators provide services that pest control should conserve whenever possible. Some botanical products could be less disruptive than certain conventional treatments under specific conditions, but being derived from plants is not a guarantee of innocuousness. Pyrethrum may also have an impact on other insects, not only

the target, and essential oils or their formulations may represent a danger. It's a matter of exposure: a spray landing directly on a bee is quite different from one applied in the absence of pollinators. New evidence suggests the need for increased scrutiny of repeated exposure and non-lethal effects such as alterations in behaviour and reproduction.

Use Within Integrated Pest Management

Botanical insecticides are most effective when used as a component of an integrated pest management program. Healthy planting stock, crop rotation, sanitation, resistant varieties and conservation of natural enemies can all diminish the demand for sprays. Scouting then informs growers if a pest population is rising to a level that warrants treatment. When a treatment is warranted, choice of treatment should be based on the life stage of the pest and the known efficacy of a product. Monitoring to follow-up tell us if damage has slowed or another intervention is warranted. This strategy makes pest control more responsive to actual field conditions and reduces unnecessary applications.

Resistance Still Matters

Insects have the ability to evolve if the same control pressure is consistently applied, whether the active substance was isolated from a plant or produced in a factory. The development of resistance may be affected by the complex nature of some botanical extracts; however, this does not mean they are immune to development of resistance. Omitting unnecessary sprays, observing labelled rates and integrating chemical control with cultural and biological control is still good advice. When multiple effective products are available, growers can schedule applications in a way that

is guided by local resistance information. To maintain the effectiveness of a botanical product, one must observe pest reactions over time rather than rely on the product's natural origin for assurance.

Safety from Mixing to Harvest

Safe use starts before a spray is applied to the crop. Concentrated preparations may cause skin or eye irritation, and the foliage may be burned if the dose is too high or if it is applied during hot weather. Growers should follow the products' directions when using them, which specify dilution rates, protective equipment and pre-harvest interval. Homemade formulations are another difficulty, as their efficacy and safety are rarely assessed. A small crop-area test may detect visible injury, but it cannot detect food safety or environmental effects. Consistent guidance and quality-assured products are critical when an edible leaf or fruit is treated close to harvest.

From Research Plot to Farm

Many plants have insecticidal activity in laboratory tests but only a handful become reliable products. Commercial production requires a long-term, renewable source of plant material, standardized extraction, stable formulations and compelling field data. In addition, farmers require affordable products that can be stored and applied with existing equipment. Tests must measure more than insect mortality: marketable yield, costs of treatment, effects on beneficial species and performance in various weather conditions are all important indicators of value. Field and stored-product pest research still provide new opportunities, but shows that scale and economics can be just as important as laboratory potency.



The Way Forward

Testing may be the next big breakthrough, rather than another plant with insecticidal properties. Researchers can increase the reliability of their findings by testing for active compounds, comparing formulations at relevant doses and reporting results across crops, seasons and locations. The potential for non-target exposure and whether products still work as part of a pest control programme should also be investigated. Farmers, extension workers and manufacturers can help to identify the traits that are most important in practice: predictable efficacy, easy-to-follow instructions and a reasonable price. This process of linking needs to research will lead to more potential plant compounds becoming valuable, sustainably-used products.

CONCLUSION

Botanical insecticides are based on a wealth of plant defence chemistry, but their use must be carefully selected. Neem, pyrethrum and essential oils illustrate the variety of effects that can be obtained, ranging from reduced feeding and interference with growth to fast-acting knock-down and repellency. New formulations might help to overcome some limitations of stability and delivery, but their benefits and risks must be tested in the field. For growers, the key questions are operational: Which pest is present, when is it most vulnerable, and what does the crop require? When integrated into integrated pest management, these questions, answered through scouting, provide a role for plant-based products in sustainable pest management.