



Rhizosphere Signaling Between Roots and Plant-Growth-Promoting Rhizobacteria (PGPR)

**K. Navya¹, Neeli Priyanka¹
Kosgi Mounika²
Puli Tanuja²**

¹Department of Agriculture
Heartfulness Academy of
Research and Innovation (HARI)
Deemed to be University
Kanha Shanti Vanam
Nadigama Telangana, India.

²Department of Horticulture
Heartfulness Academy of
Research and Innovation (HARI)
Deemed to be University
Kanha Shanti Vanam
Nadigama, Telangana, India



Open Access

*Corresponding Author
K. Navya*

Article History

Received: 21.09.2026

Revised: 26.09.2026

Accepted: 01.10.2026

This article is published under the terms of the [Creative Commons Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/).

INTRODUCTION

The rhizosphere is the region of soil directly influenced by plant roots. It is a highly active zone where roots interact continuously with bacteria, fungi and other soil microorganisms. Among these microorganisms, plant-growth-promoting rhizobacteria (PGPR) are particularly important because they can improve plant growth, nutrient availability, root development and plant resistance against several stresses.

The interaction between plant roots and PGPR is not a one-way process. Plants release different chemical compounds from their roots, while beneficial bacteria recognize these compounds and respond through movement, colonization and metabolic activities. This continuous exchange of chemical information is known as **rhizosphere signaling**.

Understanding this communication provides an important basis for developing sustainable agricultural practices based on beneficial microorganisms.

2. Root Exudates: The First Signal

Plant roots continuously release a variety of compounds into the surrounding soil. These compounds are collectively known as **root exudates**. They include sugars, amino acids, organic acids, phenolic compounds, flavonoids, vitamins and other secondary metabolites.

Root exudates serve several important functions. They provide nutrients and energy to microorganisms and influence the composition of microbial communities around the roots. Some compounds released by roots act as chemical signals that attract beneficial bacteria towards the root zone.

The quantity and composition of root exudates can change according to plant species, plant age, nutritional status and environmental conditions. Therefore, different crops can recruit different groups of microorganisms from the surrounding soil.

3. Chemotaxis: Movement of PGPR Towards Roots

Chemotaxis is the directed movement of microorganisms towards or away from specific chemical substances. PGPR can detect certain compounds released by plant roots and move towards areas where these compounds are present in higher concentrations.

This process helps beneficial bacteria locate plant roots in the soil. Bacterial cells possess specialized receptors that recognize chemical compounds and initiate appropriate movement.

The process can be summarized as:

Root exudation → Chemical recognition → Bacterial chemotaxis → Root attachment → Root colonization

Chemotaxis is therefore considered one of the important early steps in the establishment of beneficial plant–microbe interactions.

4. Root Colonization by PGPR

After reaching the root surface, PGPR must successfully establish themselves in the rhizosphere. Root colonization involves bacterial attachment, multiplication and adaptation to the conditions surrounding the root.

Many PGPR produce extracellular substances that help them attach to root surfaces. Some bacteria also form **biofilms**, which provide a stable microbial community on or near the root surface.

Successful root colonization allows PGPR to remain in close contact with plants and continuously perform beneficial functions. However, colonization depends on several factors, including bacterial strain, crop species, soil properties, temperature, moisture and competition with native microorganisms.

5. Communication Between Bacterial Cells

PGPR also communicate with one another through chemical signaling. One important mechanism is known as **quorum sensing**.

Quorum sensing allows bacterial cells to detect changes in their population density and coordinate particular activities. These activities may include biofilm formation, production of secondary metabolites, motility and interaction with plants.

Thus, communication in the rhizosphere occurs at different levels. Plants communicate with microorganisms through root-derived compounds, while bacterial cells communicate among themselves through microbial signaling molecules.

6. PGPR and Plant Growth Promotion

PGPR can promote plant growth through several direct and indirect mechanisms.

Important mechanisms include:

- Biological nitrogen fixation by suitable bacterial species.
- Solubilization and mobilization of nutrients.
- Improvement of phosphorus availability.
- Production or regulation of plant-growth-related substances.
- Enhancement of root development.
- Improvement of nutrient uptake.
- Production of siderophores.
- Suppression of harmful soil microorganisms.
- Improvement of plant tolerance to environmental stresses.

These mechanisms often operate together. The effectiveness of a PGPR strain depends on the bacterial species, crop, soil environment and prevailing climatic conditions.

7. Rhizosphere Signaling and Root Architecture

Communication between PGPR and plant roots can influence **root architecture**. Beneficial bacteria may affect root length, lateral root formation and root branching through microbial metabolites and interactions with plant hormonal pathways.

A well-developed root system can explore a larger volume of soil and improve the acquisition of water and nutrients. Therefore, PGPR-mediated changes in root architecture can indirectly contribute to better plant growth.

The interaction between roots and PGPR is consequently important not only for microbial colonization but also for the development and functioning of the root system.

8. PGPR and Plant Defense

Another important outcome of root–PGPR communication is the activation or priming of plant defense mechanisms.

Certain PGPR can stimulate **induced systemic resistance (ISR)**. In this process, beneficial microorganisms associated with the roots prepare the plant to respond more effectively to subsequent pathogen attack.

PGPR may produce microbial compounds or other elicitors that are recognized by the plant. These signals can activate defense-related pathways and increase the plant's ability to respond to harmful organisms.

This mechanism is particularly important because the interaction begins in the rhizosphere, while the resulting defense response can occur in different parts of the plant.

9. PGPR and Abiotic Stress

Rhizosphere signaling also contributes to plant responses to abiotic stresses such as drought, salinity, temperature extremes and nutrient deficiency.

Under stressful conditions, plants may modify their root exudates. These changes can influence the recruitment and activity of beneficial microorganisms. In turn, PGPR may help plants through improved nutrient acquisition, modification of root development and production of compounds associated with stress tolerance.

The effectiveness of PGPR under stress conditions depends on the compatibility between the bacterial strain and the host plant, as well as soil and environmental conditions.

10. Importance in Sustainable Agriculture

Understanding rhizosphere signaling has considerable potential for sustainable agriculture. Beneficial microorganisms can be incorporated into agricultural production systems as **biofertilizers, biostimulants and biocontrol agents**.

PGPR-based technologies may help improve nutrient-use efficiency, support plant growth and reduce dependence on excessive chemical inputs.

However, successful field application requires the selection of efficient bacterial strains, good root colonization ability, compatibility with the crop and adaptability to local soil and climatic conditions.

11. Future Prospects

Future research on rhizosphere signaling should focus on understanding the complex chemical communication between plant roots and microorganisms.

Modern approaches such as **metagenomics, transcriptomics, metabolomics and advanced microscopy** can help researchers identify the signals involved in microbial recruitment, root colonization and plant responses.

Development of crop-specific PGPR formulations and microbial consortia could further improve the practical application of beneficial microorganisms in agriculture and horticulture.

Greater understanding of plant–microbe communication may also support precision agriculture by enabling the selection of specific beneficial microorganisms according to crop, soil and environmental conditions.

CONCLUSION

The rhizosphere is a dynamic communication zone where plant roots and microorganisms continuously interact. Root exudates act as important chemical signals that influence the recruitment and behavior of PGPR. Beneficial bacteria respond through chemotaxis, root colonization, biofilm formation and various signaling mechanisms.

In return, PGPR can influence nutrient availability, root development, plant growth,

defense responses and tolerance to environmental stresses. Therefore, rhizosphere signaling represents an important biological mechanism connecting plants and beneficial microorganisms.

A better understanding of this complex communication can contribute to the development of efficient microbial technologies and more sustainable crop production systems.